



REFERENCE LIST FOR  
"INSTRUCTIONS FOR USE:  
MISOPROSTOL FOR PREVENTION  
OF POSTPARTUM HEMORRHAGE"

- Amant F, Spitz B, Timmerman D, Corremans A, Van Assche FA. Misoprostol compared with methylergometrine for the prevention of PPH: A double-blind randomized trial. BJOG 1999; 106: 1066-70.
- Caliskan E, Dilbaz B, Mutlu Meydanli M, Ozturk N, Narin MA, Haberal A. Oral misoprostol for the third stage of labor: A randomized controlled trial. Obstetrics & Gynecology 2003; 101: 921-28.
- Chandhiok N, Dhillon BS, Datey S, Mathur A, Saxena NC. Oral misoprostol for prevention of postpartum hemorrhage by paramedical workers in India. International Journal of Gynecology and Obstetrics 2006; 92: 170-75.
- Derman RJ, Kodkany BS, Goudar SS, Geller SE, Naik VA, Bellad MB, et al. Oral misoprostol in preventing postpartum haemorrhage in resource-poor communities: A randomized controlled trial. The Lancet 2006; 368: 1248-53.
- El-Refaey H, O'Brien P, Morafa W, Walder J, Rodeck C. Use of oral misoprostol in the prevention of postpartum haemorrhage. BJOG 1997; 104: 336-39.
- Gulmezoglu AM, Villar J, Ngoc NT, Piaggio G, Carroli G, Adetoro L, et al. WHO multicentre randomized controlled trial of misoprostol in the management of the third stage of labour. The Lancet 2001; 358: 689-95.
- Hofmeyr GJ, Nikodem VC, de Jager M, Gelbart BR. A randomized placebo controlled trial of oral misoprostol in the third stage of labor. BJOG 1998; 105: 971-75.
- Hofmeyr GJ, Nikodem VC, de Jager M, Drakely A. Side effects of oral misoprostol in the third stage of labour: A randomized, placebo controlled trial. South African Medical Journal 2001; 91: 432-5.
- Hoj L, Cardoso P, Nielson BB, Hvidman L, Nielson J, Aaby P. Effect of sublingual misoprostol on severe postpartum haemorrhage in a primary health centre in Guinea-Bissau: Randomized double blind clinical trial. British Medical Journal 2005; 331: 723.
- Lagenbach C. Misoprostol in preventing postpartum hemorrhage: A meta-analysis. International Journal of Gynecology and Obstetrics 2006; 92: 10-18.
- Lumbiganon P, Villar J, Piaggio G, Gulmezoglu AM, Adetoro L, Carroli G. Side effects of oral misoprostol during the first 24 hours after administration in the third stage of labor. BJOG 2002; 109: 1222-26.
- Ng PS, Chan AS, Sin WK, Tang LC, Cheung KB, Yuen PM. A multicentre randomized controlled trial of oral misoprostol and i.m. syntometrine in the management of the third stage of labor. Human Reproduction 2001; 16: 31-35.
- Oboro VO, Tabowe TO. A randomized controlled trial of misoprostol vs. oxytocin in the active management of the third stage of labor. BJOG 2002; 109: 1222-26.
- Prata N, Hamza S, Gypson R, Nada K, Vahidnia F, Potts M. Misoprostol and active management of the third stage of labor. International Journal of Gynecology and Obstetrics 2006; 94: 149-55.
- Surbek DV, Fehr PM, Hosli I, Holzgreve W. Oral misoprostol for third stage of labor: A randomized placebo-controlled trial. Obstetrics & Gynecology 1999; 94: 255-58.
- Walraven G, Blum J, Dampha Y, Sowe M, Morison L, Winikoff B, Sloan N. Misoprostol in the management of the third stage of labour in the home delivery setting in rural Gambia: A randomized controlled trial. BJOG 2005; Sept(112): 1277-83.
- WHO Recommendations on Prevention of Postpartum Hemorrhage. Geneva: WHO, 2007.