

REFERENCE LIST FOR "INSTRUCTIONS FOR USE:
ABORTION INDUCTION WITH MISOPROSTOL IN
PREGNANCIES THROUGH 9 WEEKS LMP"

Borgatta L, Mullally B, Vragovic O, Gittinger E, Chen A. Misoprostol as the primary agent for medical abortion in a low-income urban setting. *Contraception*. 2004 70:121-126.

Carbonell JL, Rodriguez J, Velazco A, Tanda R, Sanchez C, Barambio S, Chami S, Valero F, Mari J, de Vargas F, Salvador I. Oral and vaginal misoprostol 800 mcg every 8 h for early abortion. *Contraception*. 2003 Jun;67(6):457-462.

Singh K, Fong YF, Dong F. A viable alternative to surgical vacuum aspiration: repeated doses of intravaginal misoprostol over 9 hours for medical termination of pregnancies up to eight weeks. *BJOG*. 2003 Feb;110(2):175-80.

Zikopoulos KA, Papanikolaou EG, Kalantaridou SN, Tsanadis GD, Plachouras NI, Dalkalitsis NA, Paraskevaidis EA. Early pregnancy termination with vaginal misoprostol before and after 42 days gestation. *Hum Reprod*. 2002 Dec;17(12):3079-83.

Jain JK, Dutton C, Harwood B, Meckstroth KR, Mishell DR Jr. A prospective randomized, double-blinded, placebo-controlled trial comparing mifepristone and vaginal misoprostol to vaginal misoprostol alone for elective termination of early pregnancy. *Hum Reprod*. 2002 Jun;17(6):1477-82.

Tang OS, Miao BY, Lee SW, Ho PC. Pilot study on the use of repeated doses of sublingual misoprostol in termination of pregnancy up to 12 weeks gestation: efficacy and acceptability. *Hum Reprod*. 2002 Mar;17(3):654-8.

Tang OS, Ho PC. Pilot study on the use of sublingual misoprostol for medical abortion. *Contraception*. 2001 Nov;64 (5): 315-7.

Jain JK, Harwood B, Meckstroth KR, Mishell DR. Early pregnancy termination with vaginal misoprostol combined with loperamide and acetaminophen prophylaxis. *Contraception*. 2001 Apr;63(4):217-21.

Carbonell JL, Rodriguez J, Aragon S, Velazco A, Tanda R, Sanchez C, Barambio S, Chami S, Valero F. Vaginal misoprostol 1000 micrograms for early abortion. *Contraception*. 2001 Mar;63(3):131-6.

Carbonell JL, Velazco A, Varela L, Tanda R, Sanchez C, Barambio S, Chami S, Valero F, Aragon S, Mari J. Misoprostol for abortion at 9-12 weeks' gestation in adolescents. *Eur J Contracept Reprod Health Care*. 2001 Mar;6(1):39-45.

Velazco A, Varela L, Tanda R, Sanchez C, Barambio S, Chami S, Valero F, Aragon S, Mari J, Carbonell JL. Misoprostol for abortion up to 9 weeks' gestation in adolescents. *Eur J Contracept Reprod Health Care*. 2000 Dec;5(4):227-33.

Ngai SW, Tang OS, Chan YM, Ho PC. Vaginal misoprostol alone for medical abortion up to 9 weeks of gestation: efficacy and acceptability. *Hum Reprod*. 2000 May;15(5):1159-62.

Bugalho A, Mocumbi S, Faundes A, David E. Termination of pregnancies of <6 weeks gestation with a single dose of 800 microg of vaginal misoprostol. *Contraception*. 2000 Jan;61(1):47-50.

Jain JK, Meckstroth KR, Park M, Mishell DR Jr. A comparison of tamoxifen and misoprostol to misoprostol alone for early pregnancy termination. *Contraception*. 1999 Dec;60(6):353-6.

Tang OS, Wong KS, Tang LC, Ho PC. Pilot study on the use of repeated doses of misoprostol in termination of pregnancy at less than 9 weeks of gestation. *Adv Contracept*. 1999; 15(3):211-6.

Jain JK, Meckstroth KR, Mishell DR Jr. Early pregnancy termination with intravaginally administered sodium chloride solution-moistened misoprostol tablets: historical comparison with mifepristone and oral misoprostol. *Am J Obstet Gynecol*. 1999 Dec;181(6):1386-91.

Ozeren M, Bilekli C, Aydemir V, Bozkaya H. Methotrexate and misoprostol used alone or in combination for early abortion. *Contraception*. 1999 Jun;59(6):389-94.

Esteve JL, Varela L, Velazco A, Tanda R, Cabezas E, Sanchez C. Early abortion with 800 micrograms of misoprostol by the vaginal route. *Contraception*. 1999 Apr;59(4):219-25.

Carbonell JL, Varela L, Velazco A, Tanda R, Sanchez C. Vaginal misoprostol for abortion at 10-13 weeks' gestation. *Eur J Contracept Reprod Health Care*. 1999 Mar;4(1):35-40.

Carbonell Esteve JL, Varela L, Velazco A, Cabezas E, Tanda R, Sanchez C. Vaginal misoprostol for late first trimester abortion. *Contraception*. 1998 May;57(5):329-33.

Carbonell JL, Varela L, Velazco A, Fernandez C, Sanchez C. The use of misoprostol for abortion at < or = 9 weeks' gestation. *Eur J Contracept Reprod Health Care*. 1997 Sep;2(3):181-5.

Carbonell JL, Varela L, Velazco A, Fernandez C. The use of misoprostol for termination of early pregnancy. *Contraception*. 1997 Mar;55(3):165-8.

Koopersmith TB, Mishell DR Jr. The use of misoprostol for termination of early pregnancy. *Contraception*. 1996 Apr;53(4):238-42.

Bugalho A, Faundes A, Jamisse L, Usta M, Maria E, Bique C. Evaluation of the effectiveness of vaginal misoprostol to induce first trimester abortion. *Contraception*. 1996 Apr;53(4):244-6.

Creinin MD, Vittinghoff E. Methotrexate and misoprostol vs. misoprostol alone for early abortion. A randomized controlled trial. *JAMA*. 1994 Oct 19;272(15):1190-5.